

Neocate® Junior, Vanilla & Strawberry Dilution Chart

For Healthcare Professionals[§]



Contact Nutricia's Medical Affairs team for more comprehensive volume options at each concentration

Refer to can label for full instructions

1. Measure the required amount of warm or cool water into a container



2. Add prescribed amount of Neocate Junior to the water



3. Stir, or cover and shake, until the powder has dissolved

Approximate Household Measures*
Leveled and Unpacked

1 scoop [†]	7.3 g
1 tsp (5 mL)	2.5 g
1 Tbsp (15 mL)	7.6 g
¼ cup (60 mL)	30.3 g
½ cup (80 mL)	40.3 g
¾ cup (120 mL)	60.5 g
1 cup (240 mL)	121 g

Approximate Final Volume[‡]

fl oz	mL
4.7	140
8.2	245
11.7	350
18.7	560
24.6	735
32.9	990

Concentration
30 kcal/fl oz
1.00 kcal/mL
3.1 g PE/100 mL
Osmolalities:
Vanilla: 620 mOsm/kg
Strawberry: 610 mOsm/kg
~85.2% Free Water

Volume of Water		Amount of Neocate Powder [†]		
fl oz	mL	scoops	grams	kcal
4	120	4	29.2	138
7	210	7	51.1	241
10	300	10	73.0	345
16	480	16	117	551
21	630	21	153	724
28	840	29	212	999

32 kcal/fl oz
1.07 kcal/mL
3.3 g PE/100 mL
Est'd 655 mOsm/kg ||
~84.2% Free Water

fl oz	mL	scoops	grams	kcal
4.5	135	5	36.5	172
9	270	10	73.0	345
13.5	405	15	110	517
21	630	23	168	792
27	810	30	219	1034

fl oz	mL
5.4	160
10.7	320
16.1	480
24.9	750
32.1	965

34 kcal/fl oz
1.13 kcal/mL
3.6 g PE/100 mL
Est'd 695 mOsm/kg ||
~83.2% Free Water

fl oz	mL	scoops	grams	kcal
2.5	75	3	21.9	103
7.5	225	9	65.7	310
10	300	12	87.6	413
20	600	24	175	827
28	840	33	241	1137

fl oz	mL
3.0	90
9.0	270
12.0	360
24.1	725
33.6	1010

36 kcal/fl oz
1.20 kcal/mL
3.8 g PE/100 mL
Est'd 740 mOsm/kg ||
~82.2% Free Water

fl oz	mL	scoops	grams	kcal
5.5	165	7	51.1	241
9.5	285	12	87.6	413
15	450	19	139	655
22	660	28	204	965
27	810	34	248	1172

fl oz	mL
6.7	200
11.5	345
18.2	545
26.8	805
32.8	985

38 kcal/fl oz
1.27 kcal/mL
4.0 g PE/100 mL
Est'd 780 mOsm/kg ||
~81.2% Free Water

fl oz	mL	scoops	grams	kcal
9.5	285	13	94.9	448
12.5	375	17	124	586
17	510	23	168	792
20	600	27	197	930
26	780	35	256	1206

fl oz	mL
11.7	350
15.4	460
20.9	630
24.6	740
32.0	960

40 kcal/fl oz
1.33 kcal/mL
4.2 g PE/100 mL
Est'd 820 mOsm/kg ||
~80.2% Free Water

fl oz	mL	scoops	grams	kcal
5.5	165	8	58.4	276
9	270	13	94.9	448
20	600	29	212	999
26	780	38	277	1309

fl oz	mL
6.9	205
11.2	335
24.9	750
32.5	975

45 kcal/fl oz
1.50 kcal/mL
4.7 g PE/100 mL
Est'd 925 mOsm/kg ||
~77.8% Free Water

fl oz	mL	scoops	grams	kcal
3	90	5	36.5	172
6	180	10	73.0	345
12	360	20	146	689
19	570	32	234	1103
25	750	42	307	1447

fl oz	mL
3.9	115
7.7	230
15.4	460
24.5	735
32.2	965



*Household measures are based on standard US, dry household measures. All measures are level and unpacked. These values are approximations only and have been provided for convenience; results can vary significantly based on the individual, device, and method. Nutricia North America recommends using a scale for greatest accuracy.

†Dilutions are more accurate using weighed powder vs. scoops. Scoops should be level and unpacked. 1 level, unpacked scoop yields 7.3 grams powder and provides 34.5 kcal. 1 gram provides 4.72 kcal and displaces ~0.70 mL water.

‡This chart assumes 1 fl oz = 30 mL. Water volumes are exact, but final volume of prepared formula may vary slightly.

§Derived from average of 2 osmolalities at standard dilution; assumes direct linear relationship with formula concentration.

¶For patient safety, Nutricia directs caregivers who request mixing instructions at non-standard dilutions to their healthcare team for guidance.

Neocate® is a family of hypoallergenic, amino acid-based medical foods. Neocate is intended for use under medical supervision.

*HOUSEHOLD MEASURES: Clinicians can access an easy Household Measures Calculator in the Neocate Calculator at www.NutriciaLearningCenter.com in the GI/Allergy Clinician Resources section

Contact our Medical Affairs team with questions:
NutritionServices@Nutricia.com • (800)-365-7354

Neocate® Junior, Vanilla & Strawberry Dilution Chart

For Healthcare Professionals[§]



Contact Nutricia's Medical Affairs team for more comprehensive volume options at each concentration

Refer to can label for full instructions

1. Measure the required amount of warm or cool water into a container



2. Add prescribed amount of Neocate Junior to the water
3. Stir, or cover and shake, until the powder has dissolved



Approximate Household Measures*
Leveled and Unpacked

1 scoop [†]	7.3 g
1 tsp (5 mL)	2.5 g
1 Tbsp (15 mL)	7.6 g
¼ cup (60 mL)	30.3 g
½ cup (80 mL)	40.3 g
¾ cup (120 mL)	60.5 g
1 cup (240 mL)	121 g

Concentration
30 kcal/fl oz
1.00 kcal/mL
3.1 g PE/100 mL
Osmolalities:
Vanilla: 620 mOsm/kg
Strawberry: 610 mOsm/kg
~85.2% Free Water

Volume of Water		Amount of Neocate Powder [†]		
fl oz	mL	scoops	grams	kcal
4	120	4	29.2	138
7	210	7	51.1	241
10	300	10	73.0	345
16	480	16	117	551
21	630	21	153	724
28	840	29	212	999

Approximate Final Volume [‡]	
fl oz	mL
4.7	140
8.2	245
11.7	350
18.7	560
24.6	735
32.9	990

28 kcal/fl oz
0.93 kcal/mL
2.9 g PE/100 mL
Est'd 575 mOsm/kg ||
~86.2% Free Water

fl oz	mL	scoops	grams	kcal
7.5	225	7	51.1	241
10.5	315	10	73.0	345
15	450	14	102	482
21	630	20	146	689
29	870	27	197	930

fl oz	mL
8.7	260
12.2	365
17.4	520
24.4	730
33.6	1010

27 kcal/fl oz
0.90 kcal/mL
2.8 g PE/100 mL
Est'd 555 mOsm/kg ||
~86.7% Free Water

fl oz	mL	scoops	grams	kcal
5.5	165	5	36.5	172
10	300	9	65.7	310
19	570	17	124	586
21	630	19	139	655
28	840	25	183	861

fl oz	mL
6.4	190
11.5	345
21.9	655
24.2	725
32.3	970

26 kcal/fl oz
0.87 kcal/mL
2.7 g PE/100 mL
Est'd 535 mOsm/kg ||
~87.1% Free Water


fl oz	mL	scoops	grams	kcal
3.5	105	3	21.9	103
7	210	6	43.8	207
10.5	315	9	65.7	310
21	630	18	131	620
28	840	24	175	827

fl oz	mL
4.0	120
8.0	240
12.0	360
24.1	720
32.1	965

24 kcal/fl oz
0.80 kcal/mL
2.5 g PE/100 mL
Est'd 490 mOsm/kg ||
~88.1% Free Water

fl oz	mL	scoops	grams	kcal
5	150	4	29.2	138
7.5	225	6	43.8	207
10	300	8	58.4	276
20	600	16	117	551
28	840	22	161	758

fl oz	mL
5.7	170
8.5	255
11.4	340
22.7	680
31.7	950

22 kcal/fl oz
0.73 kcal/mL
2.3 g PE/100 mL
Est'd 450 mOsm/kg ||
~89.1% Free Water


fl oz	mL	scoops	grams	kcal
5.5	165	4	29.2	138
8.5	255	6	43.8	207
14	420	10	73.0	345
22	660	16	117	551
29	870	21	153	724

fl oz	mL
6.2	185
9.5	285
15.7	470
24.7	740
32.6	975

20 kcal/fl oz
0.67 kcal/mL
2.1 g PE/100 mL
Est'd 410 mOsm/kg ||
~90.1% Free Water

fl oz	mL	scoops	grams	kcal
11	330	7	51.1	241
14	420	9	65.7	310
22	660	14	102	482
30	900	19	139	655

fl oz	mL
12.2	365
15.5	465
24.4	730
33.2	995

*Household measures are based on standard US, dry household measures. All measures are level and unpacked. These values are approximations only and have been provided for convenience; results can vary significantly based on the individual, device, and method. Nutricia North America recommends using a scale for greatest accuracy.

†Dilutions are more accurate using weighed powder vs. scoops. Scoops should be level and unpacked. 1 level, unpacked scoop yields 7.3 grams powder and provides 34.5 kcal. 1 gram provides 4.72 kcal and displaces ~0.70 mL water.

‡This chart assumes 1 fl oz = 30 mL. Water volumes are exact, but final volume of prepared formula may vary slightly.

§ Derived from average of 2 osmolalities at standard dilution: assumes direct linear relationship with formula concentration.

¶ For patient safety, Nutricia directs caregivers who request mixing instructions at non-standard dilutions to their healthcare team for guidance.

Neocate® is a family of hypoallergenic, amino acid-based medical foods. Neocate is intended for use under medical supervision.

*HOUSEHOLD MEASURES: Clinicians can access an easy Household Measures Calculator in the Neocate Calculator at www.NutriciaLearningCenter.com in the GI/Allergy Clinician Resources section

Contact our Medical Affairs team with questions:
NutritionServices@Nutricia.com • (800)-365-7354

© 2018 Nutricia North America
Last Updated December 1, 2018

Neocate® Junior, Vanilla & Strawberry Dilution Chart

For Healthcare Professionals[§]

20 kcal/fl oz 0.67 kcal/mL

Est'd 410 mOsm/kg ||

2.1 g PE/100 mL

~90.1% Free Water



Based on an even starting volume of water:

These recipes start with exact water volumes and add whole scoops or level, unpacked household measures*.

They are designed to work well for parents/caregivers as well as a hospital formulary staff.

Please note that final volumes are approximate.

Volume of Water [†]		Amount of Neocate Powder [†]			Approximate Final Volume [†]	
fl oz	mL	scoops	grams	kcal	fl oz	mL
11	330	7	51.1	241	12.2	365
12.5	375	8	58.4	276	13.9	415
14	420	9	65.7	310	15.5	465
15.5	465	10	73.0	345	17.2	515
17	510	11	80.3	379	18.9	565
18.5	555	12	87.6	413	20.5	615
20	600	13	94.9	448	22.2	665
22	660	14	102	482	24.4	730
23	690	15	110	517	25.6	765
25	750	16	117	551	27.7	830
28	840	18	131	620	31.1	930
30	900	19	139	655	33.2	995
32	960	21	153	724	35.6	1065
36	1080	23	168	792	39.9	1200
40	1200	26	190	896	44.4	1335
44	1320	28	204	965	48.8	1465
50	1500	32	234	1103	55.5	1665
56	1680	36	263	1240	62.1	1865
60	1800	39	285	1344	66.6	2000
64	1920	41	299	1413	71.0	2130
70	2100	45	329	1551	77.7	2330
76	2280	49	358	1688	84.3	2530
80	2400	52	380	1792	88.9	2665

USING HOUSEHOLD MEASURES* or FULL CAN

Volume of Water [†]		Amount of Neocate Powder [†]			Approximate Final Volume [†]	
fl oz	mL	Dry measures*	grams	kcal	fl oz	mL
13	390	1/2 cup	60.5	286	14.4	430
17	510	2/3 cup	80.7	381	18.9	565
19	570	3/4 cup	90.8	428	21.1	635
26	780	1 cup	121	571	28.8	865
51	1530	2 cups	242	1142	56.6	1700
85	2550	FULL CAN	400	1888	94.4	2830



How to Prepare

Refer to can label for full preparation instructions

1. Measure the required amount of warm or cool water into a container.



2. Add the prescribed amount of Neocate Junior to the water.



3. Stir, or cover and shake, until the powder has dissolved.



Important Preparation Information

- If taken orally, Neocate Junior is best served chilled.
- Shake or stir Neocate Junior immediately before use.
- Discard formula left in cup 1 hour after feeding starts.
- If preparing in advance: refrigerate immediately at 35-40°F (2-4°C) and use within 24 hours.
- Do not freeze, boil, or use a microwave oven to prepare or warm formula.
- For oral and tube feeding; not for parenteral use.
- Tube feeding: Nutricia advises formula can be safely kept at room temperature up to 4 hours.

Storage

- Store unopened can at room temperature. Do not store at extreme temperatures.
- Once opened, reseal can and store in a cool, dry place (not a refrigerator). Use within 1 month.

© 2018 Nutricia North America.

*Household measures are based on standard US, dry household measures. All measures are level and unpacked. These values are approximations only and have been provided for convenience: results can vary significantly based on the individual, device, and method. Nutricia North America recommends using a scale for greatest accuracy.

†Dilutions are more accurate using weighed powder vs. scoops. Scoops should be level and unpacked. 1 level, unpacked scoop yields 7.3 grams powder and provides 34.5 kcal. 1 gram provides 4.72 kcal and displaces ~0.70 mL water.

‡This chart assumes 1 fl oz = 30 mL. Water volumes are exact, but final volume of prepared formula may vary slightly.

§Derived from average of 2 osmolalities at standard dilution; assumes direct linear relationship with formula concentration.

¶For patient safety, Nutricia directs caregivers who request mixing instructions at non-standard dilutions to their healthcare team for guidance.

Neocate® is a family of hypoallergenic, amino acid-based medical foods. Neocate is intended for use under medical supervision.

*HOUSEHOLD MEASURES: Clinicians can access an easy Household Measures Calculator in the Neocate Calculator at www.NutriciaLearningCenter.com in the GI/Allergy Clinician Resources section

Contact our Medical Affairs team with questions: NutritionServices@Nutricia.com • (800)-365-7354

Neocate® Junior, Vanilla & Strawberry Dilution Chart

For Healthcare Professionals[§]

22 kcal/fl oz 0.73 kcal/mL

Est'd 450 mOsm/kg ||

2.3 g PE/100 mL

~89.1% Free Water



Based on an even starting volume of water:

These recipes start with exact water volumes and add whole scoops or level, unpacked household measures*:

They are designed to work well for parents/caregivers as well as a hospital formulary staff.

Please note that final volumes are approximate.

Volume of Water [†]		Amount of Neocate Powder [†]			Approximate Final Volume [†]	
fl oz	mL	scoops	grams	kcal	fl oz	mL
5.5	165	4	29.2	138	6.2	185
7	210	5	36.5	172	7.9	235
8.5	255	6	43.8	207	9.5	285
11	330	8	58.4	276	12.4	370
12.5	375	9	65.7	310	14.0	420
14	420	10	73.0	345	15.7	470
17	510	12	87.6	413	19.0	570
18	540	13	94.9	448	20.2	605
21	630	15	110	517	23.6	705
22	660	16	117	551	24.7	740
24	720	17	124	586	26.9	805
25	750	18	131	620	28.1	840
28	840	20	146	689	31.4	940
29	870	21	153	724	32.6	975
32	960	23	168	792	35.9	1080
36	1080	26	190	896	40.4	1215
40	1200	29	212	999	44.9	1350
44	1320	32	234	1103	49.5	1485
50	1500	36	263	1240	56.1	1685
56	1680	40	292	1378	62.8	1885
60	1800	43	314	1482	67.3	2020
64	1920	46	336	1585	71.8	2155
70	2100	50	365	1723	78.5	2355

USING HOUSEHOLD MEASURES* or FULL CAN

Volume of Water [†]		Amount of Neocate Powder [†]			Approximate Final Volume [†]	
fl oz	mL	Dry measures*	grams	kcal	fl oz	mL
11.5	345	1/2 cup	60.5	286	12.9	385
15.5	465	2/3 cup	80.7	381	17.4	520
23	690	1 cup	121	571	25.8	775
35	1050	1 1/2 cups	182	857	39.2	1175
46	1380	2 cups	242	1142	51.6	1550
76	2280	FULL CAN	400	1888	85.8	2575



How to Prepare

Refer to can label for full preparation instructions

1. Measure the required amount of warm or cool water into a container.



2. Add the prescribed amount of Neocate Junior to the water.



3. Stir, or cover and shake, until the powder has dissolved.



Important Preparation Information

- If taken orally, Neocate Junior is best served chilled.
- Shake or stir Neocate Junior immediately before use.
- Discard formula left in cup 1 hour after feeding starts.
- If preparing in advance: refrigerate immediately at 35-40°F (2-4°C) and use within 24 hours.
- Do not freeze, boil, or use a microwave oven to prepare or warm formula.
- For oral and tube feeding; not for parenteral use.
- Tube feeding: Nutricia advises formula can be safely kept at room temperature up to 4 hours.

Storage

- Store unopened can at room temperature. Do not store at extreme temperatures.
- Once opened, reseal can and store in a cool, dry place (not a refrigerator). Use within 1 month.

© 2018 Nutricia North America.

*Household measures are based on standard US, dry household measures. All measures are level and unpacked. These values are approximations only and have been provided for convenience: results can vary significantly based on the individual, device, and method. Nutricia North America recommends using a scale for greatest accuracy.

†Dilutions are more accurate using weighed powder vs. scoops. Scoops should be level and unpacked. 1 level, unpacked scoop yields 7.3 grams powder and provides 34.5 kcal. 1 gram provides 4.72 kcal and displaces ~0.70 mL water.

‡This chart assumes 1 fl oz = 30 mL. Water volumes are exact, but final volume of prepared formula may vary slightly.

§ Derived from average of 2 osmolalities at standard dilution; assumes direct linear relationship with formula concentration.

¶ For patient safety, Nutricia directs caregivers who request mixing instructions at non-standard dilutions to their healthcare team for guidance.

Neocate® is a family of hypoallergenic, amino acid-based medical foods. Neocate is intended for use under medical supervision.

*HOUSEHOLD MEASURES: Clinicians can access an easy Household Measures Calculator in the Neocate Calculator at www.NutriciaLearningCenter.com in the GI/Allergy Clinician Resources section

Contact our Medical Affairs team with questions: NutritionServices@Nutricia.com • (800)-365-7354

Neocate® Junior, Vanilla & Strawberry Dilution Chart

For Healthcare Professionals[§]

24 kcal/fl oz 0.80 kcal/mL

Est'd 490 mOsm/kg ||

2.5 g PE/100 mL

~88.1% Free Water



Based on an even starting volume of water:

These recipes start with exact water volumes and add whole scoops or level, unpacked household measures*:

They are designed to work well for parents/caregivers as well as a hospital formulary staff.

Please note that final volumes are approximate.

Volume of Water [†]		Amount of Neocate Powder [†]			Approximate Final Volume [†]	
fl oz	mL	scoops	grams	kcal	fl oz	mL
2.5	75	2	14.6	69	2.8	85
5	150	4	29.2	138	5.7	170
7.5	225	6	43.8	207	8.5	255
9	270	7	51.1	241	10.2	305
10	300	8	58.4	276	11.4	340
14	420	11	80.3	379	15.9	475
15	450	12	87.6	413	17.0	510
18	540	14	102	482	20.4	610
19	570	15	110	517	21.6	645
20	600	16	117	551	22.7	680
23	690	18	131	620	26.1	780
24	720	19	139	655	27.2	815
25	750	20	146	689	28.4	850
27	810	21	153	724	30.6	915
28	840	22	161	758	31.7	950
32	960	25	183	861	36.3	1090
36	1080	28	204	965	40.8	1225
40	1200	32	234	1103	45.5	1365
44	1320	35	256	1206	50.0	1500
50	1500	40	292	1378	56.8	1705
56	1680	44	321	1516	63.5	1905
60	1800	47	343	1619	68.0	2040
64	1920	51	372	1757	72.7	2180

Approximate Household Measures*
Leveled and Unpacked

1 scoop [†]	7.3 g
1 tsp (5 mL)	2.5 g
1 Tbsp (15 mL)	7.6 g
¼ cup (60 mL)	30.3 g
½ cup (80 mL)	40.3 g
¾ cup (120 mL)	60.5 g
1 cup (240 mL)	121 g

USING HOUSEHOLD MEASURES* or FULL CAN

Volume of Water [†]		Amount of Neocate Powder [†]			Approximate Final Volume [†]	
fl oz	mL	Dry measures*	grams	kcal	fl oz	mL
7	210	1/3 cup	40.3	190	7.9	240
14	420	2/3 cup	80.7	381	15.9	475
21	630	1 cup	121	571	23.8	715
31	930	1 1/2 cups	182	857	35.2	1055
42	1260	2 cups	242	1142	47.6	1430
69	2070	FULL CAN	400	1888	78.7	2360



How to Prepare

Refer to can label for full preparation instructions

1. Measure the required amount of warm or cool water into a container.



2. Add the prescribed amount of Neocate Junior to the water.



3. Stir, or cover and shake, until the powder has dissolved.



Important Preparation Information

- If taken orally, Neocate Junior is best served chilled.
- Shake or stir Neocate Junior immediately before use.
- Discard formula left in cup 1 hour after feeding starts.
- If preparing in advance: refrigerate immediately at 35-40°F (2-4°C) and use within 24 hours.
- Do not freeze, boil, or use a microwave oven to prepare or warm formula.
- For oral and tube feeding; not for parenteral use.
- Tube feeding: Nutricia advises formula can be safely kept at room temperature up to 4 hours.

Storage

- Store unopened can at room temperature. Do not store at extreme temperatures.
- Once opened, reseal can and store in a cool, dry place (not a refrigerator). Use within 1 month.

© 2018 Nutricia North America.

*Household measures are based on standard US, dry household measures. All measures are level and unpacked. These values are approximations only and have been provided for convenience: results can vary significantly based on the individual, device, and method. Nutricia North America recommends using a scale for greatest accuracy.

†Dilutions are more accurate using weighed powder vs. scoops. Scoops should be level and unpacked. 1 level, unpacked scoop yields 7.3 grams powder and provides 34.5 kcal. 1 gram provides 4.72 kcal and displaces ~0.70 mL water.

‡This chart assumes 1 fl oz = 30 mL. Water volumes are exact, but final volume of prepared formula may vary slightly.

§ Derived from average of 2 osmolalities at standard dilution; assumes direct linear relationship with formula concentration.

¶ For patient safety, Nutricia directs caregivers who request mixing instructions at non-standard dilutions to their healthcare team for guidance.

Neocate® is a family of hypoallergenic, amino acid-based medical foods. Neocate is intended for use under medical supervision.

*HOUSEHOLD MEASURES: Clinicians can access an easy Household Measures Calculator in the Neocate Calculator at www.NutriciaLearningCenter.com in the GI/Allergy Clinician Resources section

Contact our Medical Affairs team with questions: NutritionServices@Nutricia.com • (800)-365-7354

Neocate® Junior, Vanilla & Strawberry Dilution Chart

For Healthcare Professionals[§]

26 kcal/fl oz 0.87 kcal/mL

Est'd 535 mOsm/kg ||

2.7 g PE/100 mL

~87.1% Free Water



Based on an even starting volume of water:

These recipes start with exact water volumes and add whole scoops or level, unpacked household measures*:

They are designed to work well for parents/caregivers as well as a hospital formulary staff.

Please note that final volumes are approximate.

Volume of Water [†]		Amount of Neocate Powder [†]			Approximate Final Volume [†]	
fl oz	mL	scoops	grams	kcal	fl oz	mL
3.5	105	3	21.9	103	4.0	120
7	210	6	43.8	207	8.0	240
8	240	7	51.1	241	9.2	275
10.5	315	9	65.7	310	12.0	360
14	420	12	87.6	413	16.0	480
15	450	13	94.9	448	17.2	515
16	480	14	102	482	18.4	550
21	630	18	131	620	24.1	720
22	660	19	139	655	25.2	755
23	690	20	146	689	26.4	790
24	720	21	153	724	27.6	825
28	840	24	175	827	32.1	965
29	870	25	183	861	33.3	1000
30	900	26	190	896	34.4	1035
31	930	27	197	930	35.6	1070
32	960	28	204	965	36.8	1105
34	1020	29	212	999	38.9	1170
36	1080	31	226	1068	41.3	1240
40	1200	35	256	1206	46.0	1380
44	1320	38	277	1309	50.5	1515
50	1500	43	314	1482	57.3	1720
56	1680	48	350	1654	64.2	1925
60	1800	52	380	1792	68.9	2065

Approximate Household Measures*
Leveled and Unpacked

1 scoop [†]	7.3 g
1 tsp (5 mL)	2.5 g
1 Tbsp (15 mL)	7.6 g
¼ cup (60 mL)	30.3 g
½ cup (80 mL)	40.3 g
¾ cup (120 mL)	60.5 g
1 cup (240 mL)	121 g

USING HOUSEHOLD MEASURES* or FULL CAN

Volume of Water [†]		Amount of Neocate Powder [†]			Approximate Final Volume [†]	
fl oz	mL	Dry measures*	grams	kcal	fl oz	mL
9.5	285	1/2 cup	60.5	286	10.9	325
19	570	1 cup	121	571	21.8	655
24	720	1 1/4 cups	151	714	27.5	825
38	1140	2 cups	242	1142	43.6	1310
48	1440	2 1/2 cups	303	1428	55.1	1650
63	1890	FULL CAN	400	1888	72.6	2180



How to Prepare

Refer to can label for full preparation instructions

1. Measure the required amount of warm or cool water into a container.



2. Add the prescribed amount of Neocate Junior to the water.



3. Stir, or cover and shake, until the powder has dissolved.



Important Preparation Information

- If taken orally, Neocate Junior is best served chilled.
- Shake or stir Neocate Junior immediately before use.
- Discard formula left in cup 1 hour after feeding starts.
- If preparing in advance: refrigerate immediately at 35-40°F (2-4°C) and use within 24 hours.
- Do not freeze, boil, or use a microwave oven to prepare or warm formula.
- For oral and tube feeding; not for parenteral use.
- Tube feeding: Nutricia advises formula can be safely kept at room temperature up to 4 hours.

Storage

- Store unopened can at room temperature. Do not store at extreme temperatures.
- Once opened, reseal can and store in a cool, dry place (not a refrigerator). Use within 1 month.

© 2018 Nutricia North America.

*Household measures are based on standard US, dry household measures. All measures are level and unpacked. These values are approximations only and have been provided for convenience: results can vary significantly based on the individual, device, and method. Nutricia North America recommends using a scale for greatest accuracy.

†Dilutions are more accurate using weighed powder vs. scoops. Scoops should be level and unpacked. 1 level, unpacked scoop yields 7.3 grams powder and provides 34.5 kcal. 1 gram provides 4.72 kcal and displaces ~0.70 mL water.

‡This chart assumes 1 fl oz = 30 mL. Water volumes are exact, but final volume of prepared formula may vary slightly.

§Derived from average of 2 osmolalities at standard dilution; assumes direct linear relationship with formula concentration.

¶For patient safety, Nutricia directs caregivers who request mixing instructions at non-standard dilutions to their healthcare team for guidance.

Neocate® is a family of hypoallergenic, amino acid-based medical foods. Neocate is intended for use under medical supervision.

*HOUSEHOLD MEASURES: Clinicians can access an easy Household Measures Calculator in the Neocate Calculator at www.NutriciaLearningCenter.com in the GI/Allergy Clinician Resources section

Contact our Medical Affairs team with questions: NutritionServices@Nutricia.com • (800)-365-7354

Neocate® Junior, Vanilla & Strawberry Dilution Chart

For Healthcare Professionals[§]

27 kcal/fl oz 0.90 kcal/mL

Est'd 555 mOsm/kg ||

2.8 g PE/100 mL

~86.7% Free Water



Based on an even starting volume of water:

These recipes start with exact water volumes and add whole scoops or level, unpacked household measures*.

They are designed to work well for parents/caregivers as well as a hospital formulary staff.

Please note that final volumes are approximate.

Volume of Water [†]		Amount of Neocate Powder [†]			Approximate Final Volume [†]	
fl oz	mL	scoops	grams	kcal	fl oz	mL
5.5	165	5	36.5	172	6.4	190
10	300	9	65.7	310	11.5	345
11	330	10	73.0	345	12.7	380
12	360	11	80.3	379	13.9	415
19	570	17	124	586	21.9	655
20	600	18	131	620	23.1	690
21	630	19	139	655	24.2	725
22	660	20	146	689	25.4	760
23	690	21	153	724	26.6	795
24	720	22	161	758	27.7	830
28	840	25	183	861	32.3	970
29	870	26	190	896	33.4	1005
30	900	27	197	930	34.6	1040
31	930	28	204	965	35.8	1075
32	960	29	212	999	36.9	1110
33	990	30	219	1034	38.1	1145
34	1020	31	226	1068	39.3	1180
36	1080	33	241	1137	41.6	1250
40	1200	36	263	1240	46.1	1385
44	1320	40	292	1378	50.8	1525
50	1500	45	329	1551	57.7	1730
56	1680	51	372	1757	64.7	1940
60	1800	54	394	1861	69.2	2075

USING HOUSEHOLD MEASURES* or FULL CAN

Volume of Water [†]		Amount of Neocate Powder [†]			Approximate Final Volume [†]	
fl oz	mL	Dry measures*	grams	kcal	fl oz	mL
18.5	555	1 cup	121	571	21.3	640
23	690	1 1/4 cups	151	714	26.5	795
32	960	1 3/4 cups	212	999	36.9	1110
37	1110	2 cups	242	1142	42.6	1280
46	1380	2 1/2 cups	303	1428	53.1	1590
61	1830	FULL CAN	400	1888	69.9	2100



How to Prepare

Refer to can label for full preparation instructions

1. Measure the required amount of warm or cool water into a container.



2. Add the prescribed amount of Neocate Junior to the water.



3. Stir, or cover and shake, until the powder has dissolved.



Important Preparation Information

- If taken orally, Neocate Junior is best served chilled.
- Shake or stir Neocate Junior immediately before use.
- Discard formula left in cup 1 hour after feeding starts.
- If preparing in advance: refrigerate immediately at 35-40°F (2-4°C) and use within 24 hours.
- Do not freeze, boil, or use a microwave oven to prepare or warm formula.
- For oral and tube feeding; not for parenteral use.
- Tube feeding: Nutricia advises formula can be safely kept at room temperature up to 4 hours.

Storage

- Store unopened can at room temperature. Do not store at extreme temperatures.
- Once opened, reseal can and store in a cool, dry place (not a refrigerator). Use within 1 month.

© 2018 Nutricia North America.

*Household measures are based on standard US, dry household measures. All measures are level and unpacked. These values are approximations only and have been provided for convenience: results can vary significantly based on the individual, device, and method. Nutricia North America recommends using a scale for greatest accuracy.

†Dilutions are more accurate using weighed powder vs. scoops. Scoops should be level and unpacked. 1 level, unpacked scoop yields 7.3 grams powder and provides 34.5 kcal. 1 gram provides 4.72 kcal and displaces ~0.70 mL water.

‡This chart assumes 1 fl oz = 30 mL. Water volumes are exact, but final volume of prepared formula may vary slightly.

§ Derived from average of 2 osmolalities at standard dilution; assumes direct linear relationship with formula concentration.

¶ For patient safety, Nutricia directs caregivers who request mixing instructions at non-standard dilutions to their healthcare team for guidance.

Neocate® is a family of hypoallergenic, amino acid-based medical foods. Neocate is intended for use under medical supervision.

*HOUSEHOLD MEASURES: Clinicians can access an easy Household Measures Calculator in the Neocate Calculator at www.NutriciaLearningCenter.com in the GI/Allergy Clinician Resources section

Contact our Medical Affairs team with questions: NutritionServices@Nutricia.com • (800)-365-7354

Neocate® Junior, Vanilla & Strawberry Dilution Chart

For Healthcare Professionals[§]

28 kcal/fl oz 0.93 kcal/mL

Est'd 575 mOsm/kg ||

2.9 g PE/100 mL

~86.2% Free Water



Based on an even starting volume of water:

These recipes start with exact water volumes and add whole scoops or level, unpacked household measures*.

They are designed to work well for parents/caregivers as well as a hospital formulary staff.

Please note that final volumes are approximate.

Volume of Water [†]		Amount of Neocate Powder [†]			Approximate Final Volume [‡]	
fl oz	mL	scoops	grams	kcal	fl oz	mL
7.5	225	7	51.1	241	8.7	260
8.5	255	8	58.4	276	9.9	295
9.5	285	9	65.7	310	11.0	330
10.5	315	10	73.0	345	12.2	365
15	450	14	102	482	17.4	520
16	480	15	110	517	18.6	555
17	510	16	117	551	19.7	590
18	540	17	124	586	20.9	625
19	570	18	131	620	22.1	660
20	600	19	139	655	23.2	695
21	630	20	146	689	24.4	730
22	660	21	153	724	25.6	765
29	870	27	197	930	33.6	1010
30	900	28	204	965	34.8	1045
31	930	29	212	999	35.9	1080
32	960	30	219	1034	37.1	1115
33	990	31	226	1068	38.3	1150
36	1080	34	248	1172	41.8	1255
38	1140	36	263	1240	44.1	1325
40	1200	38	277	1309	46.5	1395
44	1320	42	307	1447	51.2	1535
50	1500	47	343	1619	58.0	1740
54	1620	51	372	1757	62.7	1880

USING HOUSEHOLD MEASURES* or FULL CAN

Volume of Water [†]		Amount of Neocate Powder [†]			Approximate Final Volume [‡]	
fl oz	mL	Dry measures*	grams	kcal	fl oz	mL
13	390	3/4 cup	90.8	428	15.1	455
22	660	1 1/4 cups	151	714	25.5	765
26	780	1 1/2 cups	182	857	30.2	905
35	1050	2 cups	242	1142	40.6	1220
40	1200	2 1/4 cups	272	1285	46.4	1390
58	1740	FULL CAN	400	1888	67.4	2025



How to Prepare
Refer to can label for full preparation instructions

1. Measure the required amount of warm or cool water into a container.



2. Add the prescribed amount of Neocate Junior to the water.



3. Stir, or cover and shake, until the powder has dissolved.



Important Preparation Information

- If taken orally, Neocate Junior is best served chilled.
- Shake or stir Neocate Junior immediately before use.
- Discard formula left in cup 1 hour after feeding starts.
- If preparing in advance: refrigerate immediately at 35-40°F (2-4°C) and use within 24 hours.
- Do not freeze, boil, or use a microwave oven to prepare or warm formula.
- For oral and tube feeding; not for parenteral use.
- Tube feeding: Nutricia advises formula can be safely kept at room temperature up to 4 hours.

Storage

- Store unopened can at room temperature. Do not store at extreme temperatures.
- Once opened, reseal can and store in a cool, dry place (not a refrigerator). Use within 1 month.

© 2018 Nutricia North America.

*Household measures are based on standard US, dry household measures. All measures are level and unpacked. These values are approximations only and have been provided for convenience: results can vary significantly based on the individual, device, and method. Nutricia North America recommends using a scale for greatest accuracy.

†Dilutions are more accurate using weighed powder vs. scoops. Scoops should be level and unpacked. 1 level, unpacked scoop yields 7.3 grams powder and provides 34.5 kcal. 1 gram provides 4.72 kcal and displaces ~0.70 mL water.

‡This chart assumes 1 fl oz = 30 mL. Water volumes are exact, but final volume of prepared formula may vary slightly.

§ Derived from average of 2 osmolalities at standard dilution; assumes direct linear relationship with formula concentration.

¶ For patient safety, Nutricia directs caregivers who request mixing instructions at non-standard dilutions to their healthcare team for guidance.

Neocate® is a family of hypoallergenic, amino acid-based medical foods. Neocate is intended for use under medical supervision.

***HOUSEHOLD MEASURES:** Clinicians can access an easy Household Measures Calculator in the Neocate Calculator at www.NutriciaLearningCenter.com in the GI/Allergy Clinician Resources section

Contact our Medical Affairs team with questions:
NutritionServices@Nutricia.com • (800)-365-7354

Neocate® Junior, Vanilla & Strawberry Dilution Chart

For Healthcare Professionals[§]

30 kcal/fl oz 1.00 kcal/mL

Strawberry 610; Vanilla 620 mOsm/kg 3.1 g PE/100 mL ~85.2% Free Water



Based on an even starting volume of water:

These recipes start with exact water volumes and add whole scoops or level, unpacked household measures*.

They are designed to work well for parents/caregivers as well as a hospital formulary staff.

Please note that final volumes are approximate.

Volume of Water [†]		Amount of Neocate Powder [†]			Approximate Final Volume [‡]	
fl oz	mL	scoops	grams	kcal	fl oz	mL
1	30	1	7.3	34	1.2	35
2	60	2	14.6	69	2.3	70
3	90	3	21.9	103	3.5	105
4	120	4	29.2	138	4.7	140
7	210	7	51.1	241	8.2	245
9	270	9	65.7	310	10.5	315
10	300	10	73.0	345	11.7	350
12	360	12	87.6	413	14.0	420
14	420	14	102	482	16.4	490
16	480	16	117	551	18.7	560
18	540	18	131	620	21.1	630
21	630	21	153	724	24.6	735
24	720	25	183	861	28.3	850
26	780	27	197	930	30.6	920
28	840	29	212	999	32.9	990
30	900	31	226	1068	35.3	1060
32	960	33	241	1137	37.6	1130
34	1020	35	256	1206	40.0	1200
36	1080	37	270	1275	42.3	1270
38	1140	39	285	1344	44.6	1340
40	1200	41	299	1413	47.0	1410
44	1320	45	329	1551	51.7	1550
50	1500	51	372	1757	58.7	1760

USING HOUSEHOLD MEASURES* or FULL CAN

Volume of Water [†]		Amount of Neocate Powder [†]			Approximate Final Volume [‡]	
fl oz	mL	Dry measures*	grams	kcal	fl oz	mL
20.5	615	1 1/4 cups	151	714	24.0	720
21.5	645	1 1/3 cups	161	761	25.3	760
27	810	1 2/3 cups	202	952	31.7	950
38	1140	2 1/3 cups	282	1333	44.6	1340
43	1290	2 2/3 cups	323	1523	50.5	1515
54	1620	FULL CAN	400	1888	62.9	1890



How to Prepare

Refer to can label for full preparation instructions

1. Measure the required amount of warm or cool water into a container.



2. Add the prescribed amount of Neocate Junior to the water.



3. Stir, or cover and shake, until the powder has dissolved.



Important Preparation Information

- If taken orally, Neocate Junior is best served chilled.
- Shake or stir Neocate Junior immediately before use.
- Discard formula left in cup 1 hour after feeding starts.
- If preparing in advance: refrigerate immediately at 35-40°F (2-4°C) and use within 24 hours.
- Do not freeze, boil, or use a microwave oven to prepare or warm formula.
- For oral and tube feeding; not for parenteral use.
- Tube feeding: Nutricia advises formula can be safely kept at room temperature up to 4 hours.

Storage

- Store unopened can at room temperature. Do not store at extreme temperatures.
- Once opened, reseal can and store in a cool, dry place (not a refrigerator). Use within 1 month.

© 2018 Nutricia North America.

*Household measures are based on standard US, dry household measures. All measures are level and unpacked. These values are approximations only and have been provided for convenience: results can vary significantly based on the individual, device, and method. Nutricia North America recommends using a scale for greatest accuracy.

†Dilutions are more accurate using weighed powder vs. scoops. Scoops should be level and unpacked. 1 level, unpacked scoop yields 7.3 grams powder and provides 34.5 kcal. 1 gram provides 4.72 kcal and displaces ~0.70 mL water.

‡This chart assumes 1 fl oz = 30 mL. Water volumes are exact, but final volume of prepared formula may vary slightly.

§For patient safety, Nutricia directs caregivers who request mixing instructions at non-standard dilutions to their healthcare team for guidance. Neocate® is a family of hypoallergenic, amino acid-based medical foods. Neocate is intended for use under medical supervision.

*HOUSEHOLD MEASURES: Clinicians can access an easy Household Measures Calculator in the Neocate Calculator at www.NutriciaLearningCenter.com in the GI/Allergy Clinician Resources section

Contact our Medical Affairs team with questions: NutritionServices@Nutricia.com • (800)-365-7354

Neocate® Junior, Vanilla & Strawberry Mixing Chart

For Parents & Caregivers[§]

30 kcal/fl oz 1.00 kcal/mL



Based on an even starting volume of water:

These recipes start with exact water volumes and add whole scoops.

Please note that final volumes are approximate.

Volume of Water [†]		Amount of Neocate Powder [†]			Approximate Final Volume [‡]	
fl oz	mL	scoops	grams	kcal	fl oz	mL
1	30	1	7.3	34	1.2	35
2	60	2	14.6	69	2.3	70
3	90	3	21.9	103	3.5	105
4	120	4	29.2	138	4.7	140
5	150	5	36.5	172	5.9	175
6	180	6	43.8	207	7.0	210
7	210	7	51.1	241	8.2	245
8	240	8	58.4	276	9.4	280
9	270	9	65.7	310	10.5	315
10	300	10	73.0	345	11.7	350
11	330	11	80.3	379	12.9	385
12	360	12	87.6	413	14.0	420
13	390	13	94.9	448	15.2	455
14	420	14	102	482	16.4	490
15	450	15	110	517	17.6	525
16	480	16	117	551	18.7	560
17	510	17	124	586	19.9	595
18	540	18	131	620	21.1	630
19	570	19	139	655	22.2	665
20	600	20	146	689	23.4	700
21	630	21	153	724	24.6	735
23	690	24	175	827	27.1	815
25	750	26	190	896	29.4	885
27	810	28	204	965	31.8	955
30	900	31	226	1068	35.3	1060
33	990	34	248	1172	38.8	1165
36	1080	37	270	1275	42.3	1270
39	1170	40	292	1378	45.8	1375
42	1260	43	314	1482	49.3	1480
46	1380	47	343	1619	54.0	1620
50	1500	51	372	1757	58.7	1760
54	1620	FULL CAN	400	1888	62.9	1890



How to Prepare
Refer to can label for full preparation instructions

1. Measure the required amount of warm or cool water into a container.



2. Add the prescribed amount of **Neocate Junior** to the water.



3. Stir, or cover and shake, until the powder has dissolved.



Important Preparation Information

- If taken orally, **Neocate Junior** is best served chilled.
- Shake or stir **Neocate Junior** immediately before use.
- Discard formula left in cup 1 hour after feeding starts.
- If preparing in advance: refrigerate immediately at 35-40°F (2-4°C) and use within 24 hours.
- Do not freeze, boil, or use a microwave oven to prepare or warm formula.
- For oral and tube feeding; not for parenteral use.
- Tube feeding: Nutricia advises formula can be safely kept at room temperature up to 4 hours.

Storage

- Store unopened can at room temperature. Do not store at extreme temperatures.
- Once opened, reseal can and store in a cool, dry place (not a refrigerator). Use within 1 month.

© 2018
Nutricia North America.

Nutricia North America recommends using a scale for greatest accuracy.

The scoops Nutricia provides are validated for use with Nutricia formulas, so are preferable to household measures.

†Dilutions are more accurate using weighed powder vs. scoops. Scoops should be level and unpacked. 1 level, unpacked scoop yields 7.3 grams powder and provides 34.5 kcal. 1 gram provides 4.72 kcal and displaces ~0.70 mL water.

‡This chart assumes 1 fl oz = 30 mL. Water volumes are exact, but final volumes of prepared formula may vary slightly.

§For patient safety, Nutricia directs caregivers who request mixing instructions at non-standard dilutions, or for recipes that use household measures instead of a scale or the provided scoops, to their healthcare team for guidance.

Neocate® is a family of hypoallergenic, amino acid-based medical foods. Neocate is intended for use under medical supervision.

Contact our Nutrition Specialists with questions:
NutritionServices @ Nutricia.com
1-800-NEOCATE

Last Updated December 1, 2018

Neocate® Junior, Vanilla & Strawberry Dilution Chart

For Healthcare Professionals[§]

32 kcal/fl oz 1.07 kcal/mL

Est'd 655 mOsm/kg ||

3.3 g PE/100 mL

~84.2% Free Water



Based on an even starting volume of water:

These recipes start with exact water volumes and add whole scoops or level, unpacked household measures:^{*}

They are designed to work well for parents/caregivers as well as a hospital formulary staff.

Please note that final volumes are approximate.

Volume of Water [†]		Amount of Neocate Powder [†]			Approximate Final Volume [‡]	
fl oz	mL	scoops	grams	kcal	fl oz	mL
4.5	135	5	36.5	172	5.4	160
5.5	165	6	43.8	207	6.5	195
9	270	10	73.0	345	10.7	320
10	300	11	80.3	379	11.9	355
11	330	12	87.6	413	13.0	390
13.5	405	15	110	517	16.1	480
14.5	435	16	117	551	17.2	515
15.5	465	17	124	586	18.4	550
16.5	495	18	131	620	19.6	585
18	540	20	146	689	21.4	640
19	570	21	153	724	22.6	675
20	600	22	161	758	23.7	710
21	630	23	168	792	24.9	750
22	660	24	175	827	26.1	785
26	780	29	212	999	30.9	930
27	810	30	219	1034	32.1	965
28	840	31	226	1068	33.3	1000
30	900	33	241	1137	35.6	1070
32	960	35	256	1206	38.0	1140
36	1080	40	292	1378	42.8	1285
40	1200	44	321	1516	47.5	1425
44	1320	49	358	1688	52.3	1570
48	1440	53	387	1826	57.0	1710

USING HOUSEHOLD MEASURES* or FULL CAN

Volume of Water [†]		Amount of Neocate Powder [†]			Approximate Final Volume [‡]	
fl oz	mL	Dry measures*	grams	kcal	fl oz	mL
5	150	1/3 cup	40.3	190	5.9	180
10	300	2/3 cup	80.7	381	11.9	355
15	450	1 cup	121	571	17.8	535
30	900	2 cups	242	1142	35.6	1070
45	1350	3 cups	363	1713	53.5	1605
50	1500	FULL CAN	400	1888	59.0	1770



How to Prepare

Refer to can label for full preparation instructions

1. Measure the required amount of warm or cool water into a container.



2. Add the prescribed amount of Neocate Junior to the water.



3. Stir, or cover and shake, until the powder has dissolved.



Important Preparation Information

- If taken orally, Neocate Junior is best served chilled.
- Shake or stir Neocate Junior immediately before use.
- Discard formula left in cup 1 hour after feeding starts.
- If preparing in advance: refrigerate immediately at 35-40°F (2-4°C) and use within 24 hours.

- Do not freeze, boil, or use a microwave oven to prepare or warm formula.
- For oral and tube feeding; not for parenteral use.
- Tube feeding: Nutricia advises formula can be safely kept at room temperature up to 4 hours.

Storage

- Store unopened can at room temperature. Do not store at extreme temperatures.
- Once opened, reseal can and store in a cool, dry place (not a refrigerator). Use within 1 month.

© 2018 Nutricia North America.

*Household measures are based on standard US, dry household measures. All measures are level and unpacked. These values are approximations only and have been provided for convenience: results can vary significantly based on the individual, device, and method. Nutricia North America recommends using a scale for greatest accuracy.

†Dilutions are more accurate using weighed powder vs. scoops. Scoops should be level and unpacked. 1 level, unpacked scoop yields 7.3 grams powder and provides 34.5 kcal. 1 gram provides 4.72 kcal and displaces ~0.70 mL water.

‡This chart assumes 1 fl oz = 30 mL. Water volumes are exact, but final volume of prepared formula may vary slightly.

§ Derived from average of 2 osmolalities at standard dilution; assumes direct linear relationship with formula concentration.

¶ For patient safety, Nutricia directs caregivers who request mixing instructions at non-standard dilutions to their healthcare team for guidance.

Neocate® is a family of hypoallergenic, amino acid-based medical foods. Neocate is intended for use under medical supervision.

*HOUSEHOLD MEASURES: Clinicians can access an easy Household Measures Calculator in the Neocate Calculator at www.NutriciaLearningCenter.com in the GI/Allergy Clinician Resources section

Contact our Medical Affairs team with questions: NutritionServices@Nutricia.com • (800)-365-7354

Neocate® Junior, Vanilla & Strawberry Dilution Chart

For Healthcare Professionals[§]

34 kcal/fl oz 1.13 kcal/mL

Est'd 695 mOsm/kg ||

3.6 g PE/100 mL

~83.2% Free Water



Based on an even starting volume of water:

These recipes start with exact water volumes and add whole scoops or level, unpacked household measures*.

They are designed to work well for parents/caregivers as well as hospital formulary staff.

Please note that final volumes are approximate.

Volume of Water [†]		Amount of Neocate Powder [†]			Approximate Final Volume [‡]	
fl oz	mL	scoops	grams	kcal	fl oz	mL
2.5	75	3	21.9	103	3.0	90
5	150	6	43.8	207	6.0	180
7.5	225	9	65.7	310	9.0	270
8.5	255	10	73.0	345	10.2	305
10	300	12	87.6	413	12.0	360
11	330	13	94.9	448	13.2	395
12.5	375	15	110	517	15.1	450
13.5	405	16	117	551	16.2	485
15	450	18	131	620	18.1	540
16	480	19	139	655	19.2	575
17	510	20	146	689	20.4	610
20	600	24	175	827	24.1	725
21	630	25	183	861	25.3	760
22	660	26	190	896	26.4	795
23	690	27	197	930	27.6	830
25	750	30	219	1034	30.1	905
26	780	31	226	1068	31.3	940
28	840	33	241	1137	33.6	1010
30	900	36	263	1240	36.1	1085
32	960	38	277	1309	38.5	1155
36	1080	43	314	1482	43.3	1300
40	1200	47	343	1619	48.0	1440
44	1320	52	380	1792	52.9	1585

USING HOUSEHOLD MEASURES* or FULL CAN

Volume of Water [†]		Amount of Neocate Powder [†]			Approximate Final Volume [‡]	
fl oz	mL	Dry measures*	grams	kcal	fl oz	mL
3.5	105	1/4 cup	30.3	143	4.2	125
7	210	1/2 cup	60.5	286	8.4	250
14	420	1 cup	121	571	16.8	505
21	630	1 1/2 cups	182	857	25.2	755
28	840	2 cups	242	1142	33.6	1010
46	1380	FULL CAN	400	1888	55.5	1665



How to Prepare

Refer to can label for full preparation instructions

1. Measure the required amount of warm or cool water into a container.



2. Add the prescribed amount of Neocate Junior to the water.



3. Stir, or cover and shake, until the powder has dissolved.



Important Preparation Information

- If taken orally, Neocate Junior is best served chilled.
- Shake or stir Neocate Junior immediately before use.
- Discard formula left in cup 1 hour after feeding starts.
- If preparing in advance: refrigerate immediately at 35-40°F (2-4°C) and use within 24 hours.
- Do not freeze, boil, or use a microwave oven to prepare or warm formula.
- For oral and tube feeding; not for parenteral use.
- Tube feeding: Nutricia advises formula can be safely kept at room temperature up to 4 hours.

Storage

- Store unopened can at room temperature. Do not store at extreme temperatures.
- Once opened, reseal can and store in a cool, dry place (not a refrigerator). Use within 1 month.

© 2018 Nutricia North America.

*Household measures are based on standard US, dry household measures. All measures are level and unpacked. These values are approximations only and have been provided for convenience: results can vary significantly based on the individual, device, and method. Nutricia North America recommends using a scale for greatest accuracy.

†Dilutions are more accurate using weighed powder vs. scoops. Scoops should be level and unpacked. 1 level, unpacked scoop yields 7.3 grams powder and provides 34.5 kcal. 1 gram provides 4.72 kcal and displaces ~0.70 mL water.

‡This chart assumes 1 fl oz = 30 mL. Water volumes are exact, but final volume of prepared formula may vary slightly.

§ Derived from average of 2 osmolalities at standard dilution; assumes direct linear relationship with formula concentration.

¶ For patient safety, Nutricia directs caregivers who request mixing instructions at non-standard dilutions to their healthcare team for guidance.

Neocate® is a family of hypoallergenic, amino acid-based medical foods. Neocate is intended for use under medical supervision.

*HOUSEHOLD MEASURES: Clinicians can access an easy Household Measures Calculator in the Neocate Calculator at www.NutriciaLearningCenter.com in the GI/Allergy Clinician Resources section

Contact our Medical Affairs team with questions: NutritionServices@Nutricia.com • (800)-365-7354

Neocate® Junior, Vanilla & Strawberry Dilution Chart

For Healthcare Professionals[§]

36 kcal/fl oz 1.20 kcal/mL

Est'd 740 mOsm/kg ||

3.8 g PE/100 mL

~82.2% Free Water



Based on an even starting volume of water:

These recipes start with exact water volumes and add whole scoops or level, unpacked household measures*.

They are designed to work well for parents/caregivers as well as a hospital formulary staff.

Please note that final volumes are approximate.

Volume of Water [†]		Amount of Neocate Powder [†]			Approximate Final Volume [‡]	
fl oz	mL	scoops	grams	kcal	fl oz	mL
5.5	165	7	51.1	241	6.7	200
9.5	285	12	87.6	413	11.5	345
11	330	14	102	482	13.4	400
12.5	375	16	117	551	15.2	455
13.5	405	17	124	586	16.4	490
15	450	19	139	655	18.2	545
16.5	495	21	153	724	20.1	600
17.5	525	22	161	758	21.2	635
18	540	23	168	792	21.9	660
19	570	24	175	827	23.1	695
22	660	28	204	965	26.8	805
23	690	29	212	999	27.9	840
25	750	32	234	1103	30.5	915
26	780	33	241	1137	31.6	950
27	810	34	248	1172	32.8	985
29	870	37	270	1275	35.3	1060
30	900	38	277	1309	36.5	1095
31	930	39	285	1344	37.6	1130
32	960	41	299	1413	39.0	1170
33	990	42	307	1447	40.2	1205
35	1050	44	321	1516	42.5	1275
36	1080	46	336	1585	43.8	1315
40	1200	51	372	1757	48.7	1460

USING HOUSEHOLD MEASURES* or FULL CAN

Volume of Water [†]		Amount of Neocate Powder [†]			Approximate Final Volume [‡]	
fl oz	mL	Dry measures*	grams	kcal	fl oz	mL
6.5	195	1/2 cup	60.5	286	7.9	235
13	390	1 cup	121	571	15.8	475
23	690	1 3/4 cups	212	999	27.9	840
26	780	2 cups	242	1142	31.6	950
36	1080	2 3/4 cups	333	1571	43.8	1315
43	1290	FULL CAN	400	1888	52.4	1575



How to Prepare
Refer to can label for full preparation instructions

1. Measure the required amount of warm or cool water into a container.



2. Add the prescribed amount of Neocate Junior to the water.



3. Stir, or cover and shake, until the powder has dissolved.



Important Preparation Information

- If taken orally, Neocate Junior is best served chilled.
- Shake or stir Neocate Junior immediately before use.
- Discard formula left in cup 1 hour after feeding starts.
- If preparing in advance: refrigerate immediately at 35-40°F (2-4°C) and use within 24 hours.

- Do not freeze, boil, or use a microwave oven to prepare or warm formula.
- For oral and tube feeding; not for parenteral use.
- Tube feeding: Nutricia advises formula can be safely kept at room temperature up to 4 hours.

Storage

- Store unopened can at room temperature. Do not store at extreme temperatures.
- Once opened, reseal can and store in a cool, dry place (not a refrigerator). Use within 1 month.

© 2018 Nutricia North America.

*Household measures are based on standard US, dry household measures. All measures are level and unpacked. These values are approximations only and have been provided for convenience: results can vary significantly based on the individual, device, and method. Nutricia North America recommends using a scale for greatest accuracy.

†Dilutions are more accurate using weighed powder vs. scoops. Scoops should be level and unpacked. 1 level, unpacked scoop yields 7.3 grams powder and provides 34.5 kcal. 1 gram provides 4.72 kcal and displaces ~0.70 mL water.

‡This chart assumes 1 fl oz = 30 mL. Water volumes are exact, but final volume of prepared formula may vary slightly.

§Derived from average of 2 osmolalities at standard dilution; assumes direct linear relationship with formula concentration.

¶For patient safety, Nutricia directs caregivers who request mixing instructions at non-standard dilutions to their healthcare team for guidance.

Neocate® is a family of hypoallergenic, amino acid-based medical foods. Neocate is intended for use under medical supervision.

*HOUSEHOLD MEASURES: Clinicians can access an easy Household Measures Calculator in the Neocate Calculator at www.NutriciaLearningCenter.com in the GI/Allergy Clinician Resources section

Contact our Medical Affairs team with questions: NutritionServices@Nutricia.com • (800)-365-7354

Neocate® Junior, Vanilla & Strawberry Dilution Chart

For Healthcare Professionals[§]

38 kcal/fl oz 1.27 kcal/mL

Est'd 780 mOsm/kg ||

4.0 g PE/100 mL

~81.2% Free Water



Based on an even starting volume of water:

These recipes start with exact water volumes and add whole scoops or level, unpacked household measures*.

They are designed to work well for parents/caregivers as well as a hospital formulary staff.

Please note that final volumes are approximate.

Volume of Water [†]		Amount of Neocate Powder [†]			Approximate Final Volume [‡]	
fl oz	mL	scoops	grams	kcal	fl oz	mL
9.5	285	13	94.9	448	11.7	350
11	330	15	110	517	13.6	405
12.5	375	17	124	586	15.4	460
14	420	19	139	655	17.2	515
15.5	465	21	153	724	19.1	570
17	510	23	168	792	20.9	630
19	570	26	190	896	23.4	705
20	600	27	197	930	24.6	740
22	660	30	219	1034	27.1	815
23	690	31	226	1068	28.3	850
25	750	34	248	1172	30.8	925
26	780	35	256	1206	32.0	960
27	810	37	270	1275	33.3	1000
28	840	38	277	1309	34.5	1035
29	870	39	285	1344	35.6	1070
30	900	41	299	1413	37.0	1110
31	930	42	307	1447	38.2	1145
32	960	43	314	1482	39.3	1180
33	990	45	329	1551	40.7	1220
34	1020	46	336	1585	41.8	1255
35	1050	48	350	1654	43.2	1295
36	1080	49	358	1688	44.3	1330
38	1140	52	380	1792	46.9	1405

USING HOUSEHOLD MEASURES* or FULL CAN

Volume of Water [†]		Amount of Neocate Powder [†]			Approximate Final Volume [‡]	
fl oz	mL	Dry measures*	grams	kcal	fl oz	mL
3	90	1/4 cup	30.3	143	3.7	110
18	540	1 1/2 cups	182	857	22.2	665
24	720	2 cups	242	1142	29.6	890
31	930	2 1/2 cups	303	1428	38.1	1140
37	1110	3 cups	363	1713	45.5	1365
40	1200	FULL CAN	400	1888	49.7	1490



How to Prepare

Refer to can label for full preparation instructions

1. Measure the required amount of warm or cool water into a container.



2. Add the prescribed amount of Neocate Junior to the water.



3. Stir, or cover and shake, until the powder has dissolved.



Important Preparation Information

- If taken orally, Neocate Junior is best served chilled.
- Shake or stir Neocate Junior immediately before use.
- Discard formula left in cup 1 hour after feeding starts.
- If preparing in advance: refrigerate immediately at 35-40°F (2-4°C) and use within 24 hours.
- Do not freeze, boil, or use a microwave oven to prepare or warm formula.
- For oral and tube feeding; not for parenteral use.
- Tube feeding: Nutricia advises formula can be safely kept at room temperature up to 4 hours.

Storage

- Store unopened can at room temperature. Do not store at extreme temperatures.
- Once opened, reseal can and store in a cool, dry place (not a refrigerator). Use within 1 month.

© 2018 Nutricia North America.

*Household measures are based on standard US, dry household measures. All measures are level and unpacked. These values are approximations only and have been provided for convenience: results can vary significantly based on the individual, device, and method. Nutricia North America recommends using a scale for greatest accuracy.

†Dilutions are more accurate using weighed powder vs. scoops. Scoops should be level and unpacked. 1 level, unpacked scoop yields 7.3 grams powder and provides 34.5 kcal. 1 gram provides 4.72 kcal and displaces ~0.70 mL water.

‡This chart assumes 1 fl oz = 30 mL. Water volumes are exact, but final volume of prepared formula may vary slightly.

§ Derived from average of 2 osmolalities at standard dilution; assumes direct linear relationship with formula concentration.

¶ For patient safety, Nutricia directs caregivers who request mixing instructions at non-standard dilutions to their healthcare team for guidance.

Neocate® is a family of hypoallergenic, amino acid-based medical foods. Neocate is intended for use under medical supervision.

*HOUSEHOLD MEASURES: Clinicians can access an easy Household Measures Calculator in the Neocate Calculator at www.NutriciaLearningCenter.com in the GI/Allergy Clinician Resources section

Contact our Medical Affairs team with questions: NutritionServices@Nutricia.com • (800)-365-7354

Neocate® Junior, Vanilla & Strawberry Dilution Chart

For Healthcare Professionals[§]

40 kcal/fl oz 1.33 kcal/mL

Est'd 820 mOsm/kg ||

4.2 g PE/100 mL

~80.2% Free Water



Based on an even starting volume of water:

These recipes start with exact water volumes and add whole scoops or level, unpacked household measures*.

They are designed to work well for parents/caregivers as well as a hospital formulary staff.

Please note that final volumes are approximate.

Volume of Water [†]		Amount of Neocate Powder [†]			Approximate Final Volume [†]	
fl oz	mL	scoops	grams	kcal	fl oz	mL
5.5	165	8	58.4	276	6.9	205
9	270	13	94.9	448	11.2	335
11	330	16	117	551	13.7	410
12.5	375	18	131	620	15.6	465
13	390	19	139	655	16.2	485
16	480	23	168	792	19.9	600
18	540	26	190	896	22.4	675
20	600	29	212	999	24.9	750
22	660	32	234	1103	27.5	825
23	690	33	241	1137	28.6	860
24	720	35	256	1206	30.0	900
25	750	36	263	1240	31.1	935
26	780	38	277	1309	32.5	975
27	810	39	285	1344	33.6	1010
29	870	42	307	1447	36.2	1085
30	900	43	314	1482	37.3	1120
31	930	45	329	1551	38.7	1160
32	960	46	336	1585	39.8	1195
33	990	48	350	1654	41.2	1235
34	1020	49	358	1688	42.3	1270
35	1050	51	372	1757	43.7	1310
36	1080	52	380	1792	44.9	1345
37	1110	54	394	1861	46.2	1385

USING HOUSEHOLD MEASURES* or FULL CAN

Volume of Water [†]		Amount of Neocate Powder [†]			Approximate Final Volume [†]	
fl oz	mL	Dry measures*	grams	kcal	fl oz	mL
11.5	345	1 cup	121	571	14.3	430
19	570	1 2/3 cups	202	952	23.7	710
20	600	1 3/4 cups	212	999	24.9	750
23	690	2 cups	242	1142	28.6	860
26	780	2 1/4 cups	272	1285	32.4	970
38	1140	FULL CAN	400	1888	47.2	1415



How to Prepare
Refer to can label for full preparation instructions

1. Measure the required amount of warm or cool water into a container.



2. Add the prescribed amount of Neocate Junior to the water.



3. Stir, or cover and shake, until the powder has dissolved.



Important Preparation Information

- If taken orally, Neocate Junior is best served chilled.
- Shake or stir Neocate Junior immediately before use.
- Discard formula left in cup 1 hour after feeding starts.
- If preparing in advance: refrigerate immediately at 35-40°F (2-4°C) and use within 24 hours.
- Do not freeze, boil, or use a microwave oven to prepare or warm formula.
- For oral and tube feeding; not for parenteral use.
- Tube feeding: Nutricia advises formula can be safely kept at room temperature up to 4 hours.

Storage

- Store unopened can at room temperature. Do not store at extreme temperatures.
- Once opened, reseal can and store in a cool, dry place (not a refrigerator). Use within 1 month.

© 2018 Nutricia North America.

*Household measures are based on standard US, dry household measures. All measures are level and unpacked. These values are approximations only and have been provided for convenience: results can vary significantly based on the individual, device, and method. Nutricia North America recommends using a scale for greatest accuracy.

†Dilutions are more accurate using weighed powder vs. scoops. Scoops should be level and unpacked. 1 level, unpacked scoop yields 7.3 grams powder and provides 34.5 kcal. 1 gram provides 4.72 kcal and displaces ~0.70 mL water.

‡This chart assumes 1 fl oz = 30 mL. Water volumes are exact, but final volume of prepared formula may vary slightly.

§ Derived from average of 2 osmolalities at standard dilution; assumes direct linear relationship with formula concentration.

¶ For patient safety, Nutricia directs caregivers who request mixing instructions at non-standard dilutions to their healthcare team for guidance.

Neocate® is a family of hypoallergenic, amino acid-based medical foods. Neocate is intended for use under medical supervision.

*HOUSEHOLD MEASURES: Clinicians can access an easy Household Measures Calculator in the Neocate Calculator at www.NutriciaLearningCenter.com in the GI/Allergy Clinician Resources section

Contact our Medical Affairs team with questions: NutritionServices@Nutricia.com • (800)-365-7354

Neocate® Junior, Vanilla & Strawberry Dilution Chart

For Healthcare Professionals[§]

42 kcal/fl oz 1.40 kcal/mL

Est'd 860 mOsm/kg ||

4.4 g PE/100 mL

~79.2% Free Water



Based on an even starting volume of water:

These recipes start with exact water volumes and add whole scoops or level, unpacked household measures*.

They are designed to work well for parents/caregivers as well as a hospital formulary staff.

Please note that final volumes are approximate.

Volume of Water [†]		Amount of Neocate Powder [†]			Approximate Final Volume [†]	
fl oz	mL	scoops	grams	kcal	fl oz	mL
6.5	195	10	73.0	345	8.2	245
8.5	255	13	94.9	448	10.7	320
10.5	315	16	117	551	13.2	395
11	330	17	124	586	13.9	415
13	390	20	146	689	16.4	490
15	450	23	168	792	18.9	570
17	510	26	190	896	21.4	645
17.5	525	27	197	930	22.1	665
19	570	29	212	999	23.9	720
19.5	585	30	219	1034	24.6	740
20	600	31	226	1068	25.3	760
21	630	32	234	1103	26.5	795
22	660	34	248	1172	27.8	835
24	720	37	270	1275	30.3	910
26	780	40	292	1378	32.8	985
28	840	43	314	1482	35.3	1060
29	870	45	329	1551	36.7	1100
30	900	46	336	1585	37.8	1135
31	930	48	350	1654	39.2	1175
32	960	49	358	1688	40.3	1210
33	990	51	372	1757	41.7	1250
34	1020	52	380	1792	42.9	1285
35	1050	54	394	1861	44.2	1325

USING HOUSEHOLD MEASURES* or FULL CAN

Volume of Water [†]		Amount of Neocate Powder [†]			Approximate Final Volume [†]	
fl oz	mL	Dry measures*	grams	kcal	fl oz	mL
2	60	3 Tbsp	22.7	107	2.5	75
8	240	3/4 cup	90.8	428	10.1	305
16	480	1 1/2 cups	182	857	20.2	605
19	570	1 3/4 cups	212	999	23.9	720
24	720	2 1/4 cups	272	1285	30.4	910
36	1080	FULL CAN	400	1888	45.0	1350



How to Prepare

Refer to can label for full preparation instructions

1. Measure the required amount of warm or cool water into a container.



2. Add the prescribed amount of Neocate Junior to the water.



3. Stir, or cover and shake, until the powder has dissolved.



Important Preparation Information

- If taken orally, Neocate Junior is best served chilled.
- Shake or stir Neocate Junior immediately before use.
- Discard formula left in cup 1 hour after feeding starts.
- If preparing in advance: refrigerate immediately at 35-40°F (2-4°C) and use within 24 hours.
- Do not freeze, boil, or use a microwave oven to prepare or warm formula.
- For oral and tube feeding; not for parenteral use.
- Tube feeding: Nutricia advises formula can be safely kept at room temperature up to 4 hours.

Storage

- Store unopened can at room temperature. Do not store at extreme temperatures.
- Once opened, reseal can and store in a cool, dry place (not a refrigerator). Use within 1 month.

© 2018 Nutricia North America.

*Household measures are based on standard US, dry household measures. All measures are level and unpacked. These values are approximations only and have been provided for convenience: results can vary significantly based on the individual, device, and method. Nutricia North America recommends using a scale for greatest accuracy.

†Dilutions are more accurate using weighed powder vs. scoops. Scoops should be level and unpacked. 1 level, unpacked scoop yields 7.3 grams powder and provides 34.5 kcal. 1 gram provides 4.72 kcal and displaces ~0.70 mL water.

‡This chart assumes 1 fl oz = 30 mL. Water volumes are exact, but final volume of prepared formula may vary slightly.

§Derived from average of 2 osmolalities at standard dilution; assumes direct linear relationship with formula concentration.

¶For patient safety, Nutricia directs caregivers who request mixing instructions at non-standard dilutions to their healthcare team for guidance.

Neocate® is a family of hypoallergenic, amino acid-based medical foods. Neocate is intended for use under medical supervision.

*HOUSEHOLD MEASURES: Clinicians can access an easy Household Measures Calculator in the Neocate Calculator at www.NutriciaLearningCenter.com in the GI/Allergy Clinician Resources section

Contact our Medical Affairs team with questions: NutritionServices@Nutricia.com • (800)-365-7354

Neocate® Junior, Vanilla & Strawberry Dilution Chart

For Healthcare Professionals[§]

45 kcal/fl oz 1.50 kcal/mL

Est'd 925 mOsm/kg ||

4.7 g PE/100 mL

~77.8% Free Water



Based on an even starting volume of water:

These recipes start with exact water volumes and add whole scoops or level, unpacked household measures*.

They are designed to work well for parents/caregivers as well as a hospital formulary staff.

Please note that final volumes are approximate.

Volume of Water [†]		Amount of Neocate Powder [†]			Approximate Final Volume [‡]	
fl oz	mL	scoops	grams	kcal	fl oz	mL
3	90	5	36.5	172	3.9	115
6	180	10	73.0	345	7.7	230
6.5	195	11	80.3	379	8.4	250
12	360	20	146	689	15.4	460
12.5	375	21	153	724	16.1	480
13	390	22	161	758	16.7	500
15	450	25	183	861	19.3	580
15.5	465	26	190	896	19.9	600
16	480	27	197	930	20.6	620
18	540	30	219	1034	23.1	695
19	570	32	234	1103	24.5	735
21	630	35	256	1206	27.0	810
22	660	37	270	1275	28.3	850
23	690	39	285	1344	29.6	890
24	720	40	292	1378	30.8	925
25	750	42	307	1447	32.2	965
26	780	44	321	1516	33.5	1005
27	810	45	329	1551	34.7	1040
28	840	47	343	1619	36.0	1080
29	870	49	358	1688	37.3	1120
30	900	50	365	1723	38.5	1155
31	930	52	380	1792	39.9	1195
32	960	54	394	1861	41.2	1235

Approximate Household Measures*
Leveled and Unpacked

1 scoop [†]	7.3 g
1 tsp (5 mL)	2.5 g
1 Tbsp (15 mL)	7.6 g
¼ cup (60 mL)	30.3 g
½ cup (80 mL)	40.3 g
¾ cup (120 mL)	60.5 g
1 cup (240 mL)	121 g

USING HOUSEHOLD MEASURES* or FULL CAN

Volume of Water [†]		Amount of Neocate Powder [†]			Approximate Final Volume [‡]	
fl oz	mL	Dry measures*	grams	kcal	fl oz	mL
16.5	495	1 2/3 cups	202	952	21.2	635
22	660	2 1/4 cups	272	1285	28.4	850
23	690	2 1/3 cups	282	1333	29.6	890
24.5	735	2 1/2 cups	303	1428	31.6	945
27	810	2 3/4 cups	333	1571	34.8	1045
33	990	FULL CAN	400	1888	42.0	1260



How to Prepare
Refer to can label for full preparation instructions

1. Measure the required amount of warm or cool water into a container.



2. Add the prescribed amount of Neocate Junior to the water.



3. Stir, or cover and shake, until the powder has dissolved.



Important Preparation Information

- If taken orally, Neocate Junior is best served chilled.
- Shake or stir Neocate Junior immediately before use.
- Discard formula left in cup 1 hour after feeding starts.
- If preparing in advance: refrigerate immediately at 35-40°F (2-4°C) and use within 24 hours.
- Do not freeze, boil, or use a microwave oven to prepare or warm formula.
- For oral and tube feeding; not for parenteral use.
- Tube feeding: Nutricia advises formula can be safely kept at room temperature up to 4 hours.

Storage

- Store unopened can at room temperature. Do not store at extreme temperatures.
- Once opened, reseal can and store in a cool, dry place (not a refrigerator). Use within 1 month.

© 2018 Nutricia North America.

*Household measures are based on standard US, dry household measures. All measures are level and unpacked. These values are approximations only and have been provided for convenience; results can vary significantly based on the individual, device, and method. Nutricia North America recommends using a scale for greatest accuracy.

†Dilutions are more accurate using weighed powder vs. scoops. Scoops should be level and unpacked. 1 level, unpacked scoop yields 7.3 grams powder and provides 34.5 kcal. 1 gram provides 4.72 kcal and displaces ~0.70 mL water.

‡This chart assumes 1 fl oz = 30 mL. Water volumes are exact, but final volume of prepared formula may vary slightly.

§ Derived from average of 2 osmolalities at standard dilution; assumes direct linear relationship with formula concentration.

¶ For patient safety, Nutricia directs caregivers who request mixing instructions at non-standard dilutions to their healthcare team for guidance.

Neocate® is a family of hypoallergenic, amino acid-based medical foods. Neocate is intended for use under medical supervision.

*HOUSEHOLD MEASURES: Clinicians can access an easy Household Measures Calculator in the Neocate Calculator at www.NutriciaLearningCenter.com in the GI/Allergy Clinician Resources section

Contact our Medical Affairs team with questions:
NutritionServices@Nutricia.com • (800)-365-7354