

References Supporting Pro-Stat[®] MAX

Protein Recommendation in Critically Ill and Patients with Wounds (1.2- 2.5 g/kg body wt)

- *Hoffer, J. & Bistrain, B., Appropriate protein provision in critical illness: a systematic and narrative review. *Am J Clin Nutr.*2012;96:591-600.
- Hoffer, J. & Bistrain, B. Why Critically Ill Patients are Protein Deprived. *JPEN.*2013.37:300-309.
- *McClave, S. et al. Guidelines for the Provision and Assessment of Nutrition Support Therapy in the Adult Critically Ill Patient: Society of Critical Care Medicine (SCCM) and American Society for Parenteral and Enteral Nutrition (ASPEN). *JPEN.*2009;33:277.
- European Pressure Ulcer Advisory Panel and National Pressure Ulcer Advisory Panel. Prevention and treatment of pressure ulcers: quick reference guide. Washington DC: National Pressure Ulcer Advisory Panel; 2009. Available at: http://www.epuap.org/guidelines/Final_Quick_Prevention.pdf. Accessed 7/22/2013.
- Heidegger CP, et al. Optimization of energy provision with supplemental parenteral nutrition in critically ill patients: a randomized controlled clinical trial. *Lancet.*2013;381:385-393.
- Gerlach, A. & Murphy, C., An update on Nutrition Support in the Critically Ill. *Journal of Pharmacy Practice.*2011;24(1):70-77.

Importance of High Quality Protein & PDCAAS

- *Volpi E, et al. The response of muscle protein anabolism to combined hyperaminoacidemia and glucose-induced hyperinsulinemia is impaired in the elderly. *J Clin Endocrinol Metab.*2000;85:4481-90.
- van Loon, L & Gibala, M. Dietary Protein to Support Muscle Hypertrophy. Nestle Nutrition Institute workshop series.2011.69:79-89.
- Katsanos, C et al. Whey protein in elderly results in greater muscle protein accrual than taking its constituent EAA content. *Nutr Res.*2008;28:651-658.
- Manninen, A. Protein hydrolysates in sports nutrition. *Nutr Metab.*2009;6:38.
- Institute of Medicine of the National Academies. Dietary Reference Intakes: Protein and Amino Acids. In: Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fat, Fatty Acids, Cholesterol, Protein and Amino Acids Food and Nutrition Board, National Academy of Science, Washington, DC: National Academy Press; 2005;737.
- *Symons, TB, et al. A moderate serving of high-quality protein maximally stimulates skeletal muscle protein synthesis in young and elderly subjects. *J Am Diet Assoc.*2009;109(9):1582-6.
- *Collins, Nancy. Understanding the Protein Digestibility Corrected Amino Acid Score (PDCAAS). *Ostomy Wound Manag.*2009:8-10.
- Tome D. Criteria and markers for protein quality assessment - a review. *Brit J Nutr.*2012;108:S222-S229.
- McDonald L, et al. The Protein Book: A Complete Guide for the Coach and Athlete. 1st ed. Salt Lake City, UT: Lyle McDonald Publishing;2007.
- Castellanos, V. et al, Modular Protein Supplements and Their Application to Long-Term Care. *Nutr Clin Pract.*2006;21:485.
- Millward, J., et al. Protein quality assessment: impact of expanding understanding of protein and amino acid needs for optimal health. *Am J Clin Nutr.*2008;87(suppl):1576S- 81S.

Benefits of Whey on Muscle Protein Synthesis

- *Katsanos, C, et al. Whey protein in elderly results in greater muscle protein accrual than taking its constituent EAA content. *Nutr Res.*2008;28:651-658.
- *Pennings, B, et al. Whey protein stimulates postprandial muscle protein accretion more effectively than do casein and casein hydrolysate in older men. *Am J Clin Nutr.*2011:1-9.
- Buckley, J.D, et al, Supplementation with a whey protein hydrolysate enhances recovery of muscle force-generating capacity following eccentric exercise. *J Sci Med Sport.*2010;13:178-181.
- Manninen, A. Protein hydrolysates in sports nutrition. *Nutr Metab.*2009;6:38.
- Abrahao, V. Nourishing the dysfunctional gut and whey protein. *Curr Opin Clin Nutr.*2012;15:480-484.
- Krissansen, G., Emerging Health Properties of Whey Proteins and Their Clinical Implications. *J Am Coll Nutr.*2007;26(6):713S-723S.
- Ha, E. & Zemel, M., Functional properties of whey, whey components, and essential amino acids: mechanisms underlying health benefits for active people. *J Nutr Biochem.*2003;14:251-258.

Benefits of Protein on LBM and mortality in ICU patients

- *Weijs, PJM, et al. Optimal Protein and Energy Nutrition Decreases Mortality in Mechanically Ventilated, Critically Ill Patients: A Prospective Observational Cohort Study. *JPEN.* 2012;36:60-68.
- Baldwin, C, & Parson, TJ. Dietary advice and nutrition supplements in the management of illness-related malnutrition: a systematic review. *Clin Nutr.*2004;23:1267-1279.
- Houston, DK, et al. Dietary protein intake is associated with lean mass change in older, community-dwelling adults: the Health, Aging, and Body Composition (Health ABC) Study. *Am J Clin Nutr.*2008;87:150-5.
- Alberta, C. et al. The Relationship between nutritional intake and clinical outcomes in critically ill patients: results of an international multicenter observational study. *Intens Care Med.*2009.35:1728-1737.
- Allingstrup, M, et al. Provision of protein and energy in relation to measured requirements in intensive care patients. *Clin Nutr.*2012;31:462-468.
- Heyland, D, et al. Enhanced protein-energy provision via the enteral route in critically ill patients: a single center feasibility trial of the PEP uP protocol. *Crit Care.*2010;14:R78.

Consequences of Inadequate Intake on Poor Clinical Outcomes in Acute Care

- *Tappenden, K, et al. Critical Role of Nutrition in Improving Quality of Care: An Interdisciplinary Call to Action to Address Adult Hospital Malnutrition. *JPEN.*2013;37:482-497.
- Hoffer, J. & Bistrain, B. Why Critically Ill Patients are Protein Deprived. *JPEN.*2013.37;300-309.
- Gerlach, T., & Murphy, C. An update on Nutrition Support in the Critically Ill. *Journal of Pharmacy Practice.*2011;24:70-77.
- Barbul A. Proline Precursors to Sustain Mammalian Collagen Synthesis. *J Nutr.*2008;138:2021S-2024S.
- Stuart P et al. Use of tryptophan-fortified hydrolyzed collagen for nutritional support. *J of Dietary Supp.*2008;5(4):383-400.

Benefits of Hydrolyzed Protein

- *Tang E., et al. Ingestion of whey hydrolysate, casein, or soy protein isolate: effects on mixed muscle protein synthesis at rest and following resistance exercise in young men. *J Appl Physiol.*2009;107:987-992.
- *Koopman, R., et al. Ingestion of a protein hydrolysate is accompanied by an accelerated in vivo digestion and absorption rate when compared with its intact protein. *Am J Clin Nutr.*2009;90:1-10.
- Pennings et al. Whey protein stimulates postprandial muscle protein accretion more effectively than do casein and casein hydrolysate in older men. *Am J Clin Nutr.*2011;1-9.
- Buckley, J.D. et al, Supplementation with a whey protein hydrolysate enhances recovery of muscle force-generating capacity following eccentric exercise. *J Sci Med Sport.*2010;13:178-181.
- Manninen, A., Protein hydrolysates in sports nutrition. *Nutr Metab.*2009;6:38.
- Calbet, J. Gastric emptying, gastric secretion and enterogastrone response after administration of milk proteins or their peptide hydrolysates in humans. *Eur J Nutr.*2004; 43:127–139.
- Deglaire, A., et al. Hydrolyzed dietary casein as compared with the intact protein reduces postprandial peripheral, but not whole-body, uptake of nitrogen in humans. *Am J Clin Nutr.*2009;90:1011–22.
- Potier, M., and Tome, D. Comparison of Digestibility and Quality of Intact Proteins with Their Respective Hydrolysates. *J AOAC Int.*2008;91,NO.4.

*Key studies supporting Pro-Stat MAX.